

CS3383 Lecture 1.3: Quicksort

David Bremner

January 27, 2024



Outline


Even More Divide and Conquer

Quicksort I: partitioning

Quicksort II: random pivot

Quicksort demo

```
def quicksort(A,p,q):  
    if p>=q: return  
  
    r = partition(A,p,q,randrange(p,q))  
    quicksort(A,p,r-1)  
    quicksort(A,r+1,q)
```





Partitioning subroutine

```
PARTITION( $A, p, q$ )  $\triangleright A[p \dots q]$   
   $x \leftarrow A[p]$   $\triangleright$  pivot =  $A[p]$   
   $i \leftarrow p$   
  for  $j \leftarrow p + 1$  to  $q$   
    do if  $A[j] \leq x$   
      then  $i \leftarrow i + 1$   
          exchange  $A[i] \leftrightarrow A[j]$   
  exchange  $A[p] \leftrightarrow A[i]$   
  return  $i$ 
```


Running time
= $O(n)$ for n
elements.


Invariant:







Example of partitioning







Example of partitioning







Example of partitioning






Example of partitioning







Example of partitioning







Example of partitioning







Example of partitioning







Example of partitioning







Example of partitioning







Example of partitioning






Example of partitioning





Example of partitioning




Random pivot

- ▶ `pivot ← A[random(1..n)]`
- ▶ indicator $X_k = 1$ if we generate a $k : n - k - 1$ split, 0 otherwise
- ▶ $E[X_k] = Pr[X_k = 1] = 1/n$, assuming distinct elements.


Random pivot demo

```
def partition(A,p,q,loc):
    A[p],A[loc] = A[loc],A[p]
    i = p
    for j in range(p+1,q+1):
        if A[j] <= A[p]:
            i += 1
            A[i], A[j] = A[j], A[i]
    A[i], A[p] = A[p], A[i]
    return i
```



Analysis (continued)


$$T(n) = \begin{cases} T(0) + T(n-1) + \Theta(n) & \text{if } 0 : n-1 \text{ split,} \\ T(1) + T(n-2) + \Theta(n) & \text{if } 1 : n-2 \text{ split,} \\ \vdots & \\ T(n-1) + T(0) + \Theta(n) & \text{if } n-1 : 0 \text{ split,} \end{cases}$$
$$= \sum_{k=0}^{n-1} X_k (T(k) + T(n-k-1) + \Theta(n))$$



Calculating expectation

$$E[T(n)] = E \left[\sum_{k=0}^{n-1} X_k (T(k) + T(n-k-1) + \Theta(n)) \right]$$


Take expectations of both sides.



Calculating expectation

$$\begin{aligned} E[T(n)] &= E\left[\sum_{k=0}^{n-1} X_k (T(k) + T(n-k-1) + \Theta(n))\right] \\ &= \sum_{k=0}^{n-1} E[X_k (T(k) + T(n-k-1) + \Theta(n))] \end{aligned}$$


Linearity of expectation.



Calculating expectation

$$\begin{aligned} E[T(n)] &= E\left[\sum_{k=0}^{n-1} X_k (T(k) + T(n-k-1) + \Theta(n))\right] \\ &= \sum_{k=0}^{n-1} E[X_k (T(k) + T(n-k-1) + \Theta(n))] \\ &= \sum_{k=0}^{n-1} E[X_k] \cdot E[T(k) + T(n-k-1) + \Theta(n)] \end{aligned}$$


Independence of X_k from other random choices.



Calculating expectation

$$\begin{aligned} E[T(n)] &= E\left[\sum_{k=0}^{n-1} X_k (T(k) + T(n-k-1) + \Theta(n))\right] \\ &= \sum_{k=0}^{n-1} E[X_k (T(k) + T(n-k-1) + \Theta(n))] \\ &= \sum_{k=0}^{n-1} E[X_k] \cdot E[T(k) + T(n-k-1) + \Theta(n)] \\ &= \frac{1}{n} \sum_{k=0}^{n-1} E[T(k)] + \frac{1}{n} \sum_{k=0}^{n-1} E[T(n-k-1)] + \frac{1}{n} \sum_{k=0}^{n-1} \Theta(n) \end{aligned}$$


Linearity of expectation; $E[X_k] = 1/n$.



Calculating expectation

$$\begin{aligned} E[T(n)] &= E\left[\sum_{k=0}^{n-1} X_k (T(k) + T(n-k-1) + \Theta(n))\right] \\ &= \sum_{k=0}^{n-1} E[X_k (T(k) + T(n-k-1) + \Theta(n))] \\ &= \sum_{k=0}^{n-1} E[X_k] \cdot E[T(k) + T(n-k-1) + \Theta(n)] \\ &= \frac{1}{n} \sum_{k=0}^{n-1} E[T(k)] + \frac{1}{n} \sum_{k=0}^{n-1} E[T(n-k-1)] + \frac{1}{n} \sum_{k=0}^{n-1} \Theta(n) \\ &= \frac{2}{n} \sum_{k=1}^{n-1} E[T(k)] + \Theta(n) \end{aligned}$$

Summations have identical terms.



Hairy recurrence


$$E[T(n)] = \frac{2}{n} \sum_{k=2}^{n-1} E[T(k)] + \Theta(n)$$

(The $k = 0, 1$ terms can be absorbed in the $\Theta(n)$.)

Prove: $E[T(n)] \leq an \lg n$ for constant $a > 0$.

- Choose a large enough so that $an \lg n$ dominates $E[T(n)]$ for sufficiently small $n \geq 2$.


Use fact: $\sum_{k=2}^{n-1} k \lg k \leq \frac{1}{2} n^2 \lg n - \frac{1}{8} n^2$ (exercise).



Substitution method

$$E[T(n)] \leq \frac{2}{n} \sum_{k=2}^{n-1} ak \lg k + \Theta(n)$$


Substitute inductive hypothesis.



Substitution method

$$\begin{aligned} E[T(n)] &\leq \frac{2}{n} \sum_{k=2}^{n-1} ak \lg k + \Theta(n) \\ &\leq \frac{2a}{n} \left(\frac{1}{2} n^2 \lg n - \frac{1}{8} n^2 \right) + \Theta(n) \end{aligned}$$


Use fact.



Substitution method

$$\begin{aligned} E[T(n)] &\leq \frac{2}{n} \sum_{k=2}^{n-1} ak \lg k + \Theta(n) \\ &\leq \frac{2a}{n} \left(\frac{1}{2} n^2 \lg n - \frac{1}{8} n^2 \right) + \Theta(n) \\ &= an \lg n - \left(\frac{an}{4} - \Theta(n) \right) \end{aligned}$$

Express as *desired – residual*.



Substitution method

$$\begin{aligned} E[T(n)] &\leq \frac{2}{n} \sum_{k=2}^{n-1} ak \lg k + \Theta(n) \\ &= \frac{2a}{n} \left(\frac{1}{2} n^2 \lg n - \frac{1}{8} n^2 \right) + \Theta(n) \\ &= an \lg n - \left(\frac{an}{4} - \Theta(n) \right) \\ &\leq an \lg n, \end{aligned}$$

if a is chosen large enough so that $an/4$ dominates the $\Theta(n)$.